

Late Oligocene larger foraminifera from the Komahashi-Daini Seamount, Kyushu-Palau Ridge and their tectonic significance

MIA MOHAMMAD MOHIUDDIN¹, YUJIRO OGAWA² and KUNITERU MATSUMARU³

¹Marine Geology Department, Geological Survey of Japan, Tsukuba 305-8567 Japan; on study leave from Department of Geology and Mining, the University of Rajshahi, Rajshahi 6205 Bangladesh (e-mail: mohi@gsj.go.jp)

²Institute of Geoscience, University of Tsukuba, Tsukuba 305-8571 Japan

³Department of Geology, Faculty of Education, Saitama University Urawa, 338-8570 Japan

Received 26 August, 1999; Revised manuscript accepted 1 June 2000

Abstract. A larger foraminiferal assemblage consisting of *Miogypsinella ubaghsii* (Tan), *Spiroclypeus margaritatus* (Schlumberger) and other species is described from limestone blocks dredged at two sites on the Komahashi-Daini Seamount of the Kyushu-Palau Ridge. The fauna dates the limestone samples as Late Oligocene and is correlatable with the younger part of the Minamizaki Limestone on the Ogasawara (Bonin) Islands. These shallow-water benthic foraminifera give evidence for the shallow-water attitude of the Kyushu-Palau Ridge during the Oligocene, which has been rifted, submerged, and finally subsided to the present water depth.

Key words: Komahashi-Daini Seamount, Kyushu-Palau Ridge, larger foraminifera, Late Oligocene

Introduction

The Kyushu-Palau Ridge is an about 3,000 km long submarine ridge with a general N-S trend which divides the sea floor into the Nankai Trough on the east and the Ryukyu Trench on the west (Figure 1). On the Kyushu-Palau Ridge, a series of isolated seamounts were discovered during the 1970's (Shiki *et al.*, 1974; Shiki *et al.*, 1975). The Komahashi-Daini Seamount is located near the northern margin of this ridge. During the R/V Tansei-Maru KT94-10 Cruise, which operated July 5-12, 1994, we dredged limestone samples along with many intrusive, hypabyssal and volcanic rocks such as tonalite, andesite, tuff and pumice from the Komahashi-Daini Seamount. In this study, we describe the larger foraminifera in the limestone samples and discuss the age assignment based on the foraminiferal data and their tectonic significance.

Material

During the KT94-10 cruise, samples were dredged at two sites of the Komahashi-Daini Seamount. DG-04 site is located on the northeastern slope of the north peak, and DG-05 site on the eastern slope of the major peak (Table 1 and Figure 2). Among the rock samples, one limestone sample (DG-04-01) from the northern site and two (DG-05-01 and DG-05-02) from the southern site were studied.

The limestone samples are indurated packstone or

packstone to wackstone. All these samples are moderately hard to compact, and white to creamy white in color. They contain abundant larger and smaller benthic foraminifera, together with coral biolithite, calcareous algae and mollusks.

All of the described larger foraminiferal specimens are kept in the Geological Survey of Japan, under catalogue numbers GSJF 15418 to GSJF 15427.

Results

Thirteen foraminiferal species were identified (Figures 3–8). Dominant species are *Spiroclypeus margaritatus*, *Nephrolepidina praejaponica*, *N. angulosa*, *N. marginata*, *Eulepidina ephippioides*, *Heterostegina borneensis*, *Miogypsinella ubaghsii* and *Astrotrillina howchini*. No distinct difference in species composition was found among the three samples. This assemblage was assigned an age of Te 1-4 (Tertiary e 1-4) according to the system of Far East Letter Stages, equivalent to Late Oligocene (Hashimoto *et al.*, 1980; Hashimoto and Matsumaru, 1984; Mohiuddin, 1997). Coexistence of *M. ubaghsii* and *S. margaritatus* along with *H. borneensis*, *Eulepidina*, *Miogypsinoides* and *Spiroclypeus* is indicative of a Late Oligocene age as seen in the Melinau Limestone of Sarawak, North Borneo (Adams, 1965). Moreover, Adams and Belford (1974) suggested that the association of *S. margaritatus*, *H. borneensis* and *E. ephippioides* is indicative of the Tertiary lower e, which is believed to be equivalent to the Upper Oligocene (Chattian) of Europe.




Figure 1. Index map of dredged samples used for this study.

The *M. ubaghs* - *S. margaritatus* assemblage can be correlated with the fauna of the upper member of the Minamizaki Limestone in Chichi-Jima and Minami-Jima, Bonin Islands. *Miogypsinella boninensis* (Matsumaru, 1996) described from the Bonin Islands is thought to be a junior synonym of *Miogypsinella ubaghs* (Tan, 1936). This assemblage may be correlated with the assemblage of Te Stage limestones from 1210 to 1599 feet depth in Enewetak Atoll Drill Hole and with those from 1597.5 to 1671 feet depth in Bikini Atoll Drill Hole. The *M. ubaghs* - *S. margaritatus* assemblage is also correlated with the fauna of the Bubton Limestone, Mindoro, Philippines (Hashimoto and Matsumaru, 1984). The Te Stage is regarded as corresponding to Zone P. 21 of Blow's (1969, 1979) planktonic foraminiferal zonation.

Discussion

Konda (1975) reported larger foraminifera in limestone samples dredged from the eastern slope near a peak of the




Figure 2. Location of dredge sites KT94-10 on Kyushu-Palau Ridge. Adopted from Ohara et al. (1999). Contours in meters.

Komashashi-Daini Seamount, Kyushu-Palau Ridge and assigned to the samples an age younger than Middle Miocene based on the foraminiferal assemblage. The northern half of the Kyushu-Palau Ridge was dated around 48 Ma by Ar-Ar dating of volcaniclastic and granitic rocks (Ozima et al., 1977). A similar age was also obtained from K-Ar age of augite-orthopyroxene andesite rocks in Haha-Jima of the Bonin Islands (Kaneoka et al., 1970). These age data suggest that the Izu-Ogasawara arc was juxtaposed with the northern Kyushu-Palau Ridge before the initiation of back-arc spreading in the Shikoku Basin. Moreover, larger foraminiferal age data in this study gave a Late Oligocene age for the limestone blocks of the Komahashi-Daini

Table 1. Location of dredged samples on the Kyushu-Palau Ridge.

Sample No.	Location	Latitude	Hit bottom Longitude	Water depth (m)	Latitude	Off bottom Longitude	Water depth (m)	Dredged materials
DG-04-01	KPR, Unnamed Seamount	30°02.983'N	133°19.880'E	3800	30°02.074'N	133°18.465'E	2632	tuff, pumice and limestone
DG-05-01 and DG-05-02	KPR, Komahashi- Daini Seamount	29°53.983'N	133°22.656'E	3334	29°53.160'N	133°20.992'E	2500	tonalite, andesite and limestone

Sample No.			Larger foraminiferal species
DG-04-01	DG-05-01	DG-05-02	
X	X	X	<i>Spiroclypeus margaritatus</i> (Schlumberger)
X	X		<i>Heterostegina borneensis</i> van der Vlerk
X	X		<i>Nephrolepidina praejaponica</i> Matsumaru
X	X		<i>Nephrolepidina angulosa</i> (Provale)
X	X		<i>Nephrolepidina marginata</i> (Michelotti)
X		X	<i>Miogypsinella ubaghsii</i> (Tan)
X		X	<i>Austrotrillina howchini</i> (Schlumberger)
X			<i>Eulepidina dilatata</i> (Michelotti)
	X	X	<i>Eulepidina ephippioides</i> (Jones and Chapman)
	X		<i>Amphistegina radiata</i> (Fichtel and Moll)
X			<i>Eulepidina</i> sp.
X			<i>Ammonia</i> sp.
X		X	<i>Heterostegina</i> sp.

Figure 3. Occurrence of larger foraminiferal species in dredged samples.

Seamount, which is consistent with the oldest age of the basement rocks in the Shikoku Basin (Watts and Weissel, 1975).

The association of Late Oligocene coral-bearing limestone with benthic foraminifera of shallow-sea nature and igneous rocks recognized at the Komahashi-Daini Seamount has also been reported at DSDP Site 296, south of the seamount, at a depth of 2,920m (Figure 1). This evidence suggests that volcanogenic-calcareous sedimentary sequences of Oligocene age are rather widely distributed in the northern part of the Kyushu-Palau Ridge, including the Komahashi-Daini Seamount.

In view of the paleoenvironmental nature of the larger foraminiferal assemblage consisting of *Miogypsinella*, *Spiroclypeus*, *Austrotrillina*, *Eulepidina*, *Amphistegina* and *Heterostegina*, an environment of the shallow open ocean at the shelf edge was suggested for the deposition of limestone beds of the Komahashi-Daini Seamount, as in the case of the limestone beds of the Minamizaki Limestone, Chichi-Jima (Matsumaru, 1996). Moreover, the presence of several species of *Lepidocyclus* (*Eulepidina*) associated with pyroclastic sediments in cores 56 and 57 at DSDP Site 296 indicates a neritic environment (Ujié, 1975).

In contrast to the cases of the Komahashi-Daini Seamount and of Chichi-Jima, where the Late Oligocene sediments are exposed near the seamount surface, a drill hole at DSDP Site 296 displays a considerably continuous sequence from in situ volcanic rocks through Late Oligocene shallow-water sediments. It includes larger foraminifera and pelagic calcareous ooze, suggesting a subsidence of the Kyushu-Palau Ridge (Ujié, 1975).

It is noteworthy that the northern parts of the Kyushu-Palau Ridge and the Izu-Bonin Arc resemble each other in the timing of the cessation of volcanic activity and in the final

paleoenvironment reaching a shallow-water depth. Since Uyeda and Ben-Avraham (1972) many authors have supposed that both ridges formed a single arc at the initial stage and then were divided into two arcs owing to the spreading of the Shikoku and Parce Vella Basins. This study offers a new line of supporting evidence for this hypothesis.

Conclusion

The oldest age of the Kyushu-Palau Ridge is Late Oligocene based on larger foraminifera. The benthic foraminiferal assemblage in the limestone samples is correlated with that from the upper part of the Minamizaki Limestone exposed on the Ogasawara (Bonin) Islands of the Izu-Bonin Arc. This fact suggests that the Kyushu-Palau Ridge and the Izu-Bonin Arc initially formed a single arc. Afterward the arc may have split by a spreading of the Shikoku and Parce Vella Basins.

Systematic descriptions

Family Lepidocylinidae Scheffen, 1932

Genus *Nephrolepidina* Douvillé, 1911

Nephrolepidina praejaponica Matsumaru, 1989

Figures 6.1-6.4, 6.6, 6.7, 6.9, 6.10, 7.1, 7.6-7.9

Nephrolepidina praejaponica Matsumaru. In Matsumaru and Kimura, 1989, p. 265, 267, figs. 6.1-6.13; Matsumaru et al., 1993, p. 8, figs. 2.4, 3.6-3.8.

Material.—Thirteen specimens (GSJF 15420-1-13) including one megalospheric specimen in a vertical section (GSJF 15420-1; Figure 6.1).

Description.—Tests of megalospheric specimens, GSJF

15420-1-8, are small lenticular with diameter of 3.5 to 5.5 mm and thickness of 1.5 to 2 mm. Conical pillars are from 80 μm to 100 μm in diameter, and distributed in the central part of the test surface. The embryonic chambers are of nephrolepidine type. The protoconch is subcircular with a diameter of 240 μm . The second large chamber, the deutoerococonch embraces the protoconch and has an internal diameter of 320 μm . The ratio of the inner diameter of the deutoerococonch (II) to that of the protoconch (I) is 1.3. The outer wall of the embryonic chambers is more than 25 mm thick. The equatorial chambers of arcuate form near the periembryonic chambers change from ogival to short hexagonal near the periphery. The height of the equatorial layer near the center is about 200 μm and at the periphery less than 100 μm . The lateral chambers are rectangular in shape and are arranged in a tier of 10 to 12 layers over the center. Chambers over the central area of the test have a length of more than 160 to 200 μm , a height of 45 to 60 μm , and floors and roofs 20 to 25 μm thick.

Remarks.—The present specimen has the same features of small embryonic chambers and short hexagonal equatorial chambers in as *N. praejaponica* Matsumaru from the Lower Member of the Misaki Formation, Tosa Shimizu City, Kochi Prefecture, Shikoku (Matsumaru and Kimura, 1989) and the Early Miocene (Aquitianian) Shimizu Formation (Matsumaru et al., 1993), Shikoku Island. *Nephrolepidina praejaponica* is similar to *N. japonica* (Yabe) in overall morphology, but differs from the latter in having a small test and small embryonic chambers, primitive form of the embryonic chambers, short hexagonal equatorial chambers, rectangular lateral chambers and wavy floors and roofs.

Nephrolepidina species have been reported from Zones N. 8 and N. 9 of Blow (1969) in the Japanese mainland (Yabe, 1906; Yabe and Hanzawa, 1922; Hanzawa, 1931a, b; 1964; Matsumaru, 1967, 1971a) except the Izu Peninsula and Shikoku Island (Matsumaru, 1971a; Matsumaru and Kimura, 1989).

Nephrolepidina angulosa (Provale, 1909)

Figure 6.5

Lepidocyclina tournoueri Lemoine and R. Douvillé var. *angulosa* Provale, 1909, p. 28, pl. 3, figs. 13-15.
Lepidocyclina angulosa Provale. Rutten, 1912, p. 21, figs. 1-4.
Lepidocyclina (*Nephrolepidina*) *angulosa* Provale. Hanzawa, 1957, p. 76, 77, pl. 20, figs. 1-9, pl. 21, fig. 5, pl. 22, figs. 4, 14.
Nephrolepidina angulosa (Provale). Matsumaru, 1992, p. 259, 260, figs. 1.6, 1.7.

Material.—One megalospheric specimen in a vertical section, GSJF 15421.

Remarks.—This species is characterized by having a flat-

topped central boss with stout pillars; equatorial chambers in the mature stage are hexagonal in shape; the roof and floor of the lateral chambers are straight; and the chamber cavities are narrow and long. External appearance of the shell is similar to that of *Nephrolepidina praejaponica* Matsumaru, but it differs from the latter in possessing several conical pillars formed on the flat top of the central boss.

Family Nummulitidae de Blainville, 1827

Genus *Spiroclypeus* H. Douvillé, 1905

Spiroclypeus margaritatus (Schlumberger, 1902)

Figures 4.1, 4.2, 4.4, 4.5, 4.7, 4.9, 4.10, 5.1-5.13, 8.1

Heterostegerina margaritata Schlumberger, 1902, p. 152, 153, pl. 7, fig. 4.

Spiroclypeus orbitoideus H. Douvillé, 1905, p. 460-462, pl. 14, figs. 1-6; Tan, 1937, p. 183, 184, pl. 1, figs. 2-4, pl. 2, figs. 1-13, pl. 3, figs. 1-7; Cole, 1957a, p. 332-333, pl. 95, figs. 6-12; Matsumaru, 1976a, p. 200, pl. 1, figs. 1, 8, 10; Hashimoto, Matsumaru and Sugaya, 1981, p. 59, pl. 13, fig. 8.

Spiroclypeus leupoldi van der Vlerk, 1925, p. 14, 15, pl. 2, fig. 16; pl. 5, figs. 41, 48; Yabe and Hanzawa, 1929, p. 188, pl. 24, fig. 9; Cole, 1954, p. 577, 578, pl. 208, figs. 1-19; Hanzawa, 1957, p. 45, 46, pl. 5, figs. 7-13; Matsumaru, 1974, p. 108, pl. 15, figs. 1-4, 10, 13-15, 21-23, 28; Matsumaru, 1976a, p. 199, 200, pl. 1, figs. 4-7, 14, 15, 21, 23, 4.

Spiroclypeus yabei van der Vlerk, 1925, p. 16, pl. 2, fig. 19, pl. 5, figs. 40, 50; Tan, 1937, p. 183, pl. 1, figs. 5, 6, pl. 3, figs. 10, 11, pl. 4, figs. 8-10, text-fig. 1; Cole, 1954, p. 580-581, pl. 207, figs. 1-14, pl. 208, figs. 20-26; Cole, 1957b, p. 764, pl. 239, figs. 9-10.


Spiroclypeus tidoenganensis van der Vlerk, 1925, p. 16, 17, pl. 1, fig. 12, pl. 5, figs. 42, 47; Tan, 1937, p. 183, pl. 1, fig. 10, pl. 2, figs. 4-5, pl. 3, fig. 12, pl. 4, figs. 2-5, 19-21; Hanzawa, 1957, p. 46, 47, pl. 3, figs. 1-6, pl. 4, figs. 1, 8-10; Cole, 1957a, p. 332, pl. 95, figs. 13-15; Matsumaru, 1976a, p. 200, pl. 1, figs. 3, 9, 12, 18-20, 22, pl. 6, fig. 15; Hashimoto, Matsumaru and Sugaya, 1981, p. 60, 61, pl. 13, figs. 9, 12.

Spiroclypeus margaritata (Schlumberger). Yabe and Hanzawa, 1925, p. 627-630, pl. 2, fig. 10, pl. 3, figs. 8, 9, pl. 4, figs. 3-8, text-figs. 1-4; Krijnen, 1931, p. 89, pl. 1, figs. 1-3; Tan, 1937, p. 182, 183, pl. 2, fig. 12, pl. 3, fig. 9, pl. 4, figs. 6, 7; Hanzawa, 1940, p. 789, 790, pl. 42, figs. 3-9; Cole, 1954, p. 578-580, pl. 206, figs. 10-25, pl. 207, figs. 15, 16; Matsumaru, 1974, p. 108, pl. 15, figs. 16, 24, 26; Hashimoto and Matsumaru, 1975, p. 122, pl. 13, figs. 11, 12; Hashimoto, Matsumaru and Sugaya, 1981, p. 59, 60, pl. 13, fig. 3; Matsumaru, Myint Thein and Ogawa, 1993, p. 10, 11, figs. 2-1-9, 3-1.

Spiroclypeus margaritata (Schlumberger) var. *umbonata* Yabe and Hanzawa, 1929, p. 187, 188, pl. 124, figs. 5-8.

Spiroclypeus higginsi Cole. Hanzawa, 1957, p. 45, pl. 5, figs. 1-6, 14; Cole, 1957a, p. 332, pl. 95, figs. 1-5, pl. 109, fig. 16; Cole,

→ **Figure 4.** 1, 2, 4, 5, 7, 9, 10. *Spiroclypeus margaritatus* (Schlumberger), 1 (upper), 5, 9, 10: vertical sections, x 30, (GSJF 15418-1-4) 4, 7: oblique sections, x 30, (GSJF 15418-5-6), 2: megalospheric protoconch x 200, (GSJF 15418-7). 3. *Heterostegina* sp. vertical section, x 30. 6, 8. *Amphistegina radiata* (Fichtel and Moll), 6: vertical section, x 20, (GSJF 15427-1) 8: median section, x 20, (GSJF 15427-2) 11. *Heterostegina borneensis* van der Vlerk, vertical section, x 30, (GSJF 15419).



1957b, p. 763, 764, pl. 239, figs. 11, 12, 14; Matsumaru, 1974, p. 108, pl. 15, figs. 1, 5, 8, 12, 18, 19; Matsumaru, 1976a, p. 199, pl. 1, figs. 2, 11, 16, 17.

Spiroclypeus margaritatus (Schlumberger). Matsumaru, 1996, p. 104–108, pl. 32, figs. 1–8, pl. 33, figs. 1–9.

Material.—Twenty specimens, GSJF 15418–1–20.

Description.—Test small, inflated to lenticular, bordered by a rather thin flange, central area more than 3.5 mm in diameter and 1.5 mm in thickness. Low raised pustules distributed in umbonal portion of the test having a diameter of less than 100 µm. The megalospheric embryonic chambers consist of a spherical protoconch followed by a reniform deutoerconch. The inner diameters of protoconch (DI) and deutoerconch (DII) vary from 200 to 250 µm and 450 to 550 µm, respectively with a (DII/DI) ratio of 2.2.

Remarks.—Tan (1937) divided the species of *Spiroclypeus* into the pustulate and the reticulate group. The former group is characterized by prominent pillars on the umbonal portion of the test, the later one by the development of an external reticulation of the septa at the central part of the test. *Spiroclypeus margaritatus* belongs to the pustulate group and is characterized by large and heavy pillars, thick roofs and floors in lateral chambers, and moderate sized operculate chambers.

According to Matsumaru (1996), all the *Spiroclypeus* species reported from the West Pacific region are junior synonyms of *Spiroclypeus margaritatus* (Schlumberger). This species, known from Chichi-Jima, is restricted in occurrence to the Upper Member of the Minamizaki Limestone. It has a comparatively short stratigraphic range in Te, from the top of the *Heterostegina borneensis* Zone to the base of the *Miogypsinoides dehaartii* Zone, in the Eniwetok Atoll Drill Holes (Cole, 1957b).

Genus *Heterostegina* d'Orbigny, 1826

Heterostegina borneensis van der Vlerk, 1929

Figure 4.11

Heterostegina borneensis van der Vlerk, 1929, p. 16, figs. 6a–c, 25a–b; Cole and Bridge, 1953, p. 23, pl. 2, figs. 1–3, 5; pl. 4, figs. 16–18; Hanzawa, 1957, p. 95, pl. 26, figs. 11, 19; pl. 27, figs. 4–8; Matsumaru, 1976a, p. 199, pl. 3, figs. 17–19, 21–22; Matsumaru, 1996, p. 94–96, pl. 28, figs. 1–7.

Material.—One microspheric specimen in a vertical section, GSJF 15419.

Description.—Test small, initial part evenly lenticular with a moderately wide, thin flange on distal part. Test diameter ranges from 2.2 mm to 2.7 mm; test thickness ranges from 1.0 to 1.2 mm; thickness of pillars varies from 120 µm at umbo to 100 µm at tip of flange. In vertical section, embryonic apparatus biloculine; initial protoconch subcircular; its diameter less than 100 µm. Prominent pillars are present on the central boss of the test. Pillars penetrating to outer

wall of embryonic apparatus and equatorial layer.

Remarks.—*Heterostegina borneensis* and *Spiroclypeus margaritatus* co-occur in the Lower and Upper members of the Minamizaki Limestone. In the Komahashi-Daini Seamount Limestone, *H. borneensis* is associated with *Spiroclypeus margaritatus*, the latter species being the more abundant one. *H. borneensis* has also been recognized as a marker species to distinguish Te1–4 from Te5 (Cole, 1957a; Adams, 1965; Matsumaru, 1974, 1978), since van der Vlerk (1925) regarded it to be a useful species for delimiting Te1–4.

Family Austrotrillinidae Loeblich and Tappan, 1986

Genus *Austrotrillina* Parr, 1942

Austrotrillina howchini (Schlumberger, 1893)

Figure 8.11

Trillina howchini Schlumberger, 1893, p. 119, 120, text-figs. 1–2, pl. 3, fig. 6; Hanzawa, 1940, p. 791–793, pl. 42, figs. 1, 2.

Austrotrillina howchini (Schlumberger). Cole and Bridge, 1953, p. 20, pl. 14, fig. 12; Cole, 1954, p. 573, pl. 210, figs. 6–9; Hanzawa, 1957, p. 38, pl. 22, figs. 12, 13; pl. 34, figs. 1, 2; Matsumaru, 1996, p. 214–216, pl. 84, figs. 3–7.

Material.—One microspheric specimen in a longitudinal section, GSJF 15424.

Remarks.—*Austrotrillina howchini* originally described from Saipan is also found in the Bikini Atoll Drill Holes associated with *Spiroclypeus* and *Eulepidina* in Te Stage (Cole, 1954). The stratigraphic range of this species has been given as Te through Tf1–2 (Glaessner, 1943) and as Te and Tf1 (van der Vlerk, 1948). Hanzawa (1940) stated that this species is found only in the Aquitanian stage in the Western Pacific. Hashimoto and Matsumaru (1984) suggested that *A. howchini* ranged from Te4 to Te5–Tf1. This species occurs in association with *Miogypsinella boninensis* and *Spiroclypeus margaritatus* in the Minamizaki Limestone, Chichi-Jima, assigned to Te 1–4 of the Far East Letter Stages (Hashimoto et al., 1980; Hashimoto and Matsumaru, 1984).

Family Lepidocylinidae Scheffen, 1932

Subfamily Eulepidininae Matsumaru, 1991

Genus *Eulepidina* H. Douillé, 1911

Eulepidina ephippioides (Jones and Chapman, 1900)


Figures 6.8, 7.3, 7.4


Orbitoides (Lepidocyclus) ephippioides Jones and Chapman, 1900, p. 251, 252, pl. 20, fig. 9. *Lepidocyclus ephippioides* Jones and Chapman. Grimsdale, 1952, p. 240–244, pl. 23, figs. 8, 17, 18.

Lepidocyclus (Eulepidina) formosa Schlumberger. Cole, 1954, p. 594–597, pl. 216, figs. 1–16; pl. 217, figs. 9–11, pl. 218, figs. 1, 3, 4.

Lepidocyclus (Eulepidina) gibbosa Yabe. Cole, 1954, p. 597, pl. 217, figs. 9–11.

→ **Figure 5.** 1–13. *Spiroclypeus margaritatus* (Schlumberger). 1–3, 5–10, 12, 13: vertical sections, x 30, (GSJF 15418–8–18), 4, 11: oblique sections, x 30, (GSJF 15418–19–20).





Lepidocyclina (Eulepidina) planata Oppenoorth. Cole, 1954, p. 597, 598, pl. 217, figs. 7, 8; pl. 218, figs. 5, 6.

Lepidocyclina (Eulepidina) ephippioides Jones and Chapman. Cole, 1957b, p. 346–337, pl. 108, figs. 4–13; pl. 109, figs. 11–15.

Eulepidina ephippioides (Jones and Chapman). Matsumaru, 1996, p. 178–181, pl. 65, figs. 1–6, pl. 66, figs. 1–3; pl. 67, figs. 1–6; pl. 68, figs. 1–3; pl. 69, figs. 1–4; pl. 70, figs. 1–5, text-fig. 20–5.

Material. — Three megalospheric specimens (GSJF 15426–1–3).

Remarks. — *Eulepidina ephippioides* is characterized by the possession of a small nucleoconch and hexagonal or spatulate equatorial chambers. The earliest name of this species was thought to be *Orbitoides (Lepidocyclina) ephippioides* Jones and Chapman. According to Grimsdale (1952), the American Oligocene species *L. (E.) favosa* Cushman should be a synonym of *L. ephippioides* (Jones and Chapman).

Eulepidina dilatata (Michelotti, 1861)

Figure 8.1 (lower)

Orbitoides dilatata Michelotti, 1861, p. 17, pl. 1, figs. 1–2.

Eulepidina dilatata (Michelotti). Matsumaru, 1971b, p. 184, 185, pl. 22, figs. 28–38; Hashimoto and Matsumaru, 1975, p. 114, 115, pl. 12, figs. 10, 11; Matsumaru, 1996, p. 162–178, pl. 60, figs. 1–6; pl. 61, figs. 1–6; pl. 62, figs. 1–7; pl. 63, figs. 1–6; pl. 64, figs. 1–2, text-figs. 20–2, 4, text-fig. 30.

Material. — One obliquely sectioned megalospheric specimen, GSJF 15425.

Remarks. — The present species is characterized by having a lenticular shape, polygonal outline, large nucleoconch, hexagonal equatorial chambers, low and long lateral chambers and thin roofs and floors. It differs in general shell shape from *Eulepidina ephippioides* (Jones and Chapman). Recently, Matsumaru (1996) investigated the size of the embryonic chambers of *E. dilatata* and *E. ephippioides* from the Minamizaki Limestone, Chichi-Jima and concluded that microspheric *E. dilatata* slightly differs in chamber budding formation from microspheric *E. ephippioides*.

Family Miogypsinidae Vaughan, 1928

Genus *Miogypsinella* Hanzawa, 1940

Miogypsinella ubaghsii (Tan, 1936)

Figures 7.2, 8.2, 8.3

Miogypsinoides ubaghsii Tan, 1936, p. 47, 48, pl. 1, figs. 1–7; Cole, 1954, p. 603, 604, pl. 221, figs. 5, 9–18; pl. 222, figs. 13, 15.

Miogypsinella ubaghsii (Tan). Hanzawa, 1940, p. 767, 768, text-fig. 4.

Material. — Three megalospheric specimens; one in an

equatorial section, GSJF 15423–3 (Figure 8.3), one in an axial section, GSJF 15423–1 (Figure 8.2), and one in a vertical section, GSJF 15423–2 (Figure 8.2).

Description. — Test small, slightly wider than long, fan-shaped; 1.5 to 1.8 mm in diameter and 0.65 to 0.75 mm in thickness. Surface ornamentation consists of large pustules over the initial portion and finer, closer-spaced pustules over the distal portion. Embryonic chambers are bilocular, first chamber is nearly spherical and second chamber is reniform. Initial chambers are followed by subquadrate periembryonic chambers arranged so that they form virtually two coils. Periembryonic chambers gradually increase in length as they are added for about 1.5 volutions at which point they decrease gradually in length to the end of the coil.

Remarks. — The present species differs from *Miogypsinella borodinensis* Matsumaru, 1996, described from Minamizaki Limestone, Chichi-Jima, in having fewer equatorial and embryonic chambers and a small apical angle.

Family Amphisteginidae Cushman, 1927

Genus *Amphistegina* d'Orbigny, 1826

Amphistegina radiata (Fichtel and Moll, 1798)

Figures 4.6, 4.8, 8.1

Nautilus radiatus Fichtel and Moll, 1798, p. 58, pl. 8, figs. 8a–d.

Amphistegina lessoni d'Orbigny. Yabe and Hanzawa, 1925, p. 48, 49, pl. 8, figs. 9, 10; Hanzawa, 1931b, p. 156, pl. 24, fig. 7; pl. 25, figs. 5–8; pl. 10, fig. 4.

Amphistegina radiata (Fichtel and Moll). Yabe and Hanzawa, 1929, p. 179, 180, pl. 18, fig. 6; Matsumaru, 1976b, p. 408, pl. 1, figs. 1–3, 5–13, 17, 23, 26–27, text-figs. 6–8. Matsumaru, 1996, p. 188, pl. 74, figs. 1–5.

Material. — Three microspheric specimens (GSJF 15427–1–3)

Remarks. — The present specimens show a close similarity with those of *A. radiata* described from the Minamizaki Limestones (Matsumaru, 1996) and are characterized by many chambers in the last whorl, curvature of the spiral suture and septa and a large protoconch.


Acknowledgments

We acknowledge the help of the crew and scientific party on board the R/V *Tansei Maru* KT94–10 cruise, particularly T. Ishii. Thanks are also due to H. Ujié, Takushoku University, for discussion on some larger foraminifera taxonomy and for critical reading of the manuscript. We would like to thank A. Nishimura, Geological Survey of Japan, for extensive and fruitful discussions regarding geologic history of the Kyushu-Palau Ridge.

References


Adams, C.G., 1965: The foraminifera and stratigraphy of the

◀ Figure 6. 1–4, 6, 7, 9, 10. *Nephrolepidina praejaponica* Matsumaru. vertical sections, x 30, (GSJF 15420–1–8). 5 *Nephrolepidina angulosa* (Provale), vertical section, x 30, (GSJF 15421). 8. *Eulepidina ephippioides* (Jones and Chapman), vertical section, x 20, (GSJF 15426–1).



- Melinau Limestone, Sarawak, and its importance in Tertiary correlation. *Quarterly Journal of the Geological Society of London*, vol. 121, p. 283-338, pls. 21-30.
- Adams, C.G. and Belford, D.J., 1974: Foraminiferal biostratigraphy of the Oligocene-Miocene limestones of the Christmas Island (Indian Ocean). *Palaeontology*, vol. 17, p. 475-506.
- Blainville, H.M.D. de, 1827: *Manuel de malacologie et de conchyliologie* (1825). Paris: F.G. Levraut, 664pp., 87pls.
- Blow, W.H., 1969: Late Middle Eocene to Recent planktonic foraminiferal biostratigraphy. In, Brönnemann, P. and Renz, H. H., eds., *Proceedings of the First International Conference on Planktonic Microfossils*, Geneva, 1967, vol. 1, p. 199-421. E. J. Brill, Leiden.
- Blow, W.H., 1979: *The Cainozoic Globigerinida*, 3 vols. 1413pp., E.J. Brill, Leiden.
- Cole, W.S., 1954: Larger foraminifera and smaller diagnostic foraminifera from the Bikini Drill Holes. *U. S. Geological Survey Professional Papers*, 260-O, p. 569-608.
- Cole, W.S., 1957a: Larger foraminifera of Saipan. *U.S. Geological Survey Professional Papers*, 280-I, p. 321-360.
- Cole, W.S., 1957b: Larger foraminifera from Eniwetok Atoll Drill Holes. *U.S. Geological Survey Professional Papers*, 260-V, p. 743-784.
- Cole, W.S. and Bridge, J., 1953: Geology and larger foraminifera of Saipan Island. *U.S. Geological Survey Professional Papers*, 253, p. 1-45, pls. 2-15.
- Cushman, J.A., 1927: An outline of a reclassification of foraminifera. *Contributions from the Cushman Laboratory for Foraminiferal Research*, vol. 3, 105 pp., 25 pls.
- Douville, H., 1905: Les foraminifères dans le Tertiaire de Borneo. *Bulletin de la Société Géologique de France*, Série 4, vol. 6, no. 2, p. 435-464.
- Douville, H., 1911: Les foraminifères dans le Tertiaire des Philippines. *Philippine Journal of Science*, series D, vol. 6, p. 53-80, pl. A-D.
- Fichtel, L. and Moll, J.P.C., 1798: *Testacea microscopica, aliaque minuta ex generibus Argonauta et Nautilus, ad naturam picta et descripta*. Vienna: Camesina, xii+123 pp., 1-24 pls.
- Glaessner, M.F., 1943: Problem of stratigraphic correlation in the Indo-Pacific region. *Royal Society Victoria Proceedings, New Series*, vol. 55, pt. 1, p. 41-80.
- Grimsdale, T.F., 1952: Cretaceous and Tertiary foraminifera from the Middle East. *British Museum (Natural History) Bulletin*, vol. 1, no. 8, p. 223-247, pls. 20-25.
- Hanzawa, S., 1931a: Notes on Tertiary foraminiferous rocks from the Kwanto Mountainland, Japan. *Science Reports of the Tohoku Imperial University, Second Series (Geology)*, vol. 12, no. 2, p. 141-157.
- Hanzawa, S., 1931b: On some Miocene rocks with *Lepidocyclina* from the Izu and Boso Peninsulas. *Science Reports of the Tohoku University, Second Series (Geology)*, vol. 12, p. 157-170, pls. 27, 28.
- Hanzawa, S., 1940: Micropaleontological studies of drill cores from a deep well in the Kita-Daito-Zima (North Borodino Island). *Jubilee Publication of Prof. H. Yabe's 60th Birthday*, p. 755-802 pls. 39-42.
- Hanzawa, S., 1957: Cenozoic foraminifera of Micronesia. *Geological Society of America Memoir*, 66, p. 1-63.
- Hanzawa, S., 1964: The phylomorphogenesis of the Tertiary foraminiferal families, *Lepidocyclinidae* and *Miogypsinae*. *Science Reports of the Tohoku University, 2nd Series (Geology)*, vol. 35, p. 295-313.
- Hashimoto, W. and Matsumaru, K., 1975: Larger foraminifera from the Philippines, Part III. Limestone from eastern coastal ranges of north and central Luzon. *Geology and Paleontology of Southeast Asia*, vol. 16, p. 117-125, pl. 13.
- Hashimoto, W. and Matsumaru, K., 1984: Mesozoic and Cenozoic larger foraminifera of the Philippine and a reference to those found from Borneo by the APRSA's paleontological reconnaissance. *Geology and Paleontology of Southeast Asia*, vol. 25, p. 147-166.
- Hashimoto, W., Matsumaru, K. and Fuchimoto, H., 1980: Consideration on the stratigraphy of the Caraballo Range, northern Luzon: Larger foraminiferal ranges on the Cenozoic of the Philippines. *Proceedings of First International Congress on Pacific Neogene Stratigraphy*, vol. 20, p. 119-134.
- Hashimoto, W., Matsumaru, K. and Sugaya, M., 1981: Larger foraminifera from the Philippines. Part XI. On the Coal Harbor Limestone, Cagraray Island, Batan Island Group, Albay Province. *Geology and Paleontology of Southeast Asia*, vol. 22, p. 55-62, pl. 13.
- Jones, T.R. and Chapman, F., 1900: On the foraminifera of the Orbitoidal limestones and reef rocks of Christmas Island. In, Andrews, C.W. ed., *A Monograph of Christmas Island (Indian Ocean)*, British Museum (Natural History), London, p. 226-264. pls. 20, 21.
- Kaneoka, I., Isshiki, N. and Zashu, S., 1970: K-Ar ages of the Izu-Bonin Islands. *Geochemical Journal*, vol. 4, p. 53-60.
- Konda, I., 1975: Some paleontological results and problematic subjects on GDP Research Cruise. *Marine Science/Monthly*, vol. 7, no. 7, p. 465-470. (in Japanese with English abstract)
- Krijnen, W.F., 1931: Het Genus *Spiroclypeus* in het Indo-Pacificische Gebied. *Geologisch-Mijnbouwkundig Genootschap Nederlanden Kolonien, Verhandelingen, Geology Series*, vol. 9, p. 77-111.
- Loeblich, A.R.Jr. and Tappan, H., 1986: Some new and redefined genera and families of *Textulariina*, *Fusulinina*, *Involutinina* and *Miliolina* (Foraminiferida). *Journal of Foraminiferal Research*, vol. 16, p. 334-346.
- Matsumaru, K., 1967: Geology of the Tomioka area, Gunma Prefecture, with a note on "Lepidocyclina" from the Abuta Limestone Member. *Science Reports of the Tohoku University, Second Series (Geology)*, vol. 39, p. 113-147.
- Matsumaru, K., 1971a: Studies of the Genus *Nephrolepidina* in Japan. *Science Reports of the Tohoku University, Second Series (Geology)*, vol. 42, p. 97-185, pls. 9-26.
- Matsumaru, K., 1971b: The genera *Lepidocyclina* and *Eulepidina* from New Zealand. *Transactions and*

← Figure 7. 1, 6-9. *Nephrolepidina praejaponica* Matsumaru, 1: oblique section, x 20, 6-9: vertical sections, x 30, (GSJF 15420-9—13). 2. *Miogypsinella ubaghsii* (Tan), vertical section, x 80, (GSJF 15423-1). 3, 4. *Eulepidina ephippioides* (Jones and Chapman), vertical sections, x 20, (GSJF 15426-2—3). 5. *Eulepidina* sp., vertical section, x 10.



- Proceedings of the Paleontological Society of Japan, New Series*, no. 84, p. 179–189, pls. 22, 23.
- Matsumaru, K., 1974: Larger foraminifera from east Mindanao, the Philippines. *Geology and Paleontology of Southeast Asia*, vol. 14, p. 101–115, pls. 14–19.
- Matsumaru, K., 1976a: Larger foraminifera from the islands of Saipan and Guam, Micronesia. In, Takayanagi, Y. and Saito, T. eds., *Progress in Micropaleontology*, p. 190–213. Micropaleontology Press, New York.
- Matsumaru, K., 1976b: Larger foraminifera from the Ryukyu Group, Nansei Shoto Islands, Japan. *First International Symposium on Benthonic Foraminiferal Margins. Pt. B. Maritime Sediment, Special Publication 1*, p. 401–424.
- Matsumaru, K., 1978: Biostratigraphy and paleoecological transition of larger foraminifera, Minamizaki Limestone, Chichi-Jima, Japan. *Proceedings of Second Working Group Meeting Biostratigraphic Datumplane of the Pacific Neogene, IGCP Project 114*, p. 63–88, Bandung.
- Matsumaru, K., 1991: On the evolutionary classification of the Family Lepidocyclinidae (Foraminiferida). *Transactions and Proceedings of the Paleontological Society of Japan, New Series*, no. 164, p. 883–909.
- Matsumaru, K., 1992: Some Miocene Nephrolepidina (Family Lepidocyclinidae) from the Shimoshiroiwa Formation, Izu Peninsula, Japan. In, Ishizaki, K. and Saito, T. eds., *Centenary of Japanese Micropaleontology*, p. 257–265.
- Matsumaru, K., 1996: Tertiary larger foraminifera (Foraminiferida) from the Ogasawara Islands, Japan. *Paleontological Society of Japan, Special Papers*, no. 36, 239p.
- Matsumaru, K. and Kimura, K., 1989: Larger foraminifera from the Eocene Shumizu and Miocene Misaki Formations in Tosa Shimizu City, Kochi Prefecture, Shikoku, Japan. *Transactions and Proceedings of the Paleontological Society of Japan, New Series*, no. 156, p. 156–169.
- Matsumaru, K., Myint Thein and Ogawa, Y., 1993: Early Miocene (Aquitian) larger foraminifera from the Shimizu Formation, Ashizuri Cape, Kochi Prefecture, Shikoku, Japan. *Transactions and Proceedings of the Paleontological Society of Japan, New Series*, no. 169, p. 1–14.
- Michelotti, G., 1861: Études sur le Miocene inférieur de l'Italie septentrionale. *Natuurkundige Verhandelingen Hollandsche Maatschappij der Wetenschappen*, vol. 2, pt. 15, p. 1–183, pls. 1–16.
- Mohiuddin, M.M., 1997: Biostratigraphic and tectonic significance of the Paleogene to Early Miocene carbonate rocks: Mineoka Tectonic Belt and Kyushu-Palau Ridge. Unpublished Ph. D. Thesis, University of Tsukuba, 160p., Japan.
- Ohara, Y., Kasuga, S., Kato, Y., Okino, K., Taira, A., Arima, M., Haraguchi, S., Ishii, T. and Katsura T., 1999: Continental crust formation in an oceanic island arc: drilling proposal at the Kyushu-Palau Ridge. *Chikyu Monthly, Special No. 23*, p. 133–140. (in Japanese)
- Orbigny, A.d', 1826: Tableau méthodique de la classe des Céphalopodes. *Annales des Sciences Naturelles*, vol. 7, p. 96–314, pls. 10–17.
- Ozima, M., Kaneoka, Y. and Ujiie, H., 1977: ^{40}Ar – ^{39}Ar age of rocks and development mode of the Philippine Sea. *Nature*, vol. 267, p. 816–818.
- Parr, W.J., 1942: New genera of Foraminifera from the Tertiary of Victoria. *Mining and Geological Journal*, vol. 2, p. 361–363, 5 figs.
- Provale, I., 1909: Di alcune Nummulitine e Orbitoidine dell' Isola di Borneo. *Rivista Italiana de Paleontologia*, vol. 15, p. 65–96, pls 2–3.
- Rutten, L.M.R., 1912: Studien über foraminiferen aus Ost-Asien. *Sammlungen des geologischen Reichsmuseum in Leiden, Folge 1*, vol. 9, p. 202–224, pls. 2, 13.
- Scheffen, W., 1932: Zur Morphologie und Morphogenese der "Lepidocyclinen". *Paläontologische Zeitschrift*, vol. 14, p. 233–256, pls. 9–10.
- Schlumberger, C., 1893: Note sur les genres Trillina et Linderina. *Bulletin de la Société Géologique de France, Série 3*, vol. 21, p. 118–123, pl. 3.
- Schlumberger, C., 1902: Note sur un Lepidocyclina nouveau de Borneo. *Sammlungen des Geologischen Reichsmuseums in Leiden, Folge 1*, vol. 6, p. 250–253, pl. 7.
- Shiki, T., Aoki, H., Suzuki, M., Musashino, M. and Okuda, Y., 1974: Geological and petrographical results of the GDP 8th cruises in the Philippine Sea. *Marine Sciences/Monthly*, no. 6, p. 555–560. (in Japanese)
- Shiki, T., Tokuoka, H., Aoki, H., Misawa, Y., Konda, I. and Nishida, S., 1975: Some geological results of the GDP cruises in the Philippine Sea, with special references to bottom sampling of the GDP-8, 11. In, *Geological Problems of the Philippine Sea, Geological Society of Japan*, p. 67–74.
- Tan, S.H., 1936: Zur Kenntnis der Miogypsiniden. *De Ingenieur in Nederlandsche-Indië, Afd. IV, Mijnbouw en Geologie*, Jaarg. 3, no. 3, p. 45–61, pls. 1–2.
- Tan, S.H., 1937: On the genus *Spiroclypeus* H. Douvillé with a description of Eocene *Spiroclypeus vermicularis* nov. sp. from Koetai in east Borneo. *De Ingenieur in Nederlandsche-Indië, Afd. IV, Mijnbouw en Geologie*, Jaarg. 4, no. 10, p. 177–193.
- Ujiie, H., 1975: Planktonic foraminiferal biostratigraphy in the Western Philippine Sea. In, Karig, D.E., Ingle, J.C. et al., eds. *Initial Reports of the Deep Sea Drilling Project*, vol. 31, p. 677–691. U.S. Goverment Printing Office, Washington, D.C.
- Uyeda, S. and Ben-Avraham, Z., 1972: Origin and development of the Philippine Sea. *Nature*, vol. 240, p. 176.
- Vaughan, T.W., 1928: Subfamily Miogypsinidae Vaughan. In, Cushman, J.A., 1928, *Foraminifera their classification and economic use, Special Publication of Cushman Laboratory for Foraminiferal Research*, vol. 1, 401p.
- Vlerk, I.M. van der, 1925: A study of Tertiary foraminifera from the "Tidoeengsche Landen" (E. Borneo). *Dutch East Indies, Dienst van der Mijnbouw, Wetenschappelijke Mededelingen. Dienst van den Mijnbouw in Nederland*

Figure 8. 1. Bioclastic packstone containing diagnostic species such as *Spiroclypeus margaritatus* (Schlumberger) (GSJF 15418–20), *Nephrolepidina marginata* (Michelotti) (GSJF 15422), *Eulepidina dilatata* (Michelotti) (GSJF 15425) and *Amphistegina radiata* (Fichtel and Moll) (GSJF 15427–3) x 20. 2, 3. *Miogypsinella ubaghsii* (Tan). 2: axial section, x 20, GSJF 15423–2, 3: equatorial section, x 20, (GSJF 15423–3). 4–10, 12, 13. *Ammonia* sp., 4, 7, 9, 10, 12: oblique sections, x 20, 5: equatorial section, x 20, 6, 8, 13: axial sections, x 20. 11. *Austrotrillina howchini* (Schlumberger), longitudinal section, x 20, (GSJF 15424).

- sch-Indie*, no. 3, p. 13–32, pls. 1–6.
- Vlerk, I.M. van der, 1929: Groote foraminiferen van N.O. Borneo. *Wetenschappelijke Mededelingen Dienst van der Mijnbouw in Nederlandsch-Indie* no. 9, p. 5–30.
- Vlerk, I.M. van der, 1948: Stratigraphy of the Cenozoic of the East Indies based on Foraminifera. *International Geological Congress, Report of 18th Session, Great Britain*, pt. 15, p. 61–63.
- Watts, A. B. and Weissel, J.K., 1975: Tectonic history of the Shikoku marginal basin. *Earth and Planetary Science Letters*, vol. 25, p. 239–250.
- Yabe, H., 1906: On the orbitoid limestone from Nakakosaka and from Kuboi on the Lake Kawaguchi. *Journal of Geological Society of Japan*, vol. 13, no. 156, p. 317–320.
- (in Japanese)
- Yabe, H. and Hanzawa, S., 1922: *Lepidocyclina* from Nakakosaka, Province of Kodzuke, Japan. *Japanese Journal of Geology and Geography*, vol. 1, no. 1 p. 45–50, pls. 5–8.
- Yabe, H. and Hanzawa, S., 1925: A *Lepidocyclina*-Limestone from Klias Peninsula, B. N. Borneo. *Gedenkboek Verbeek, Verhandelingen Geologisch-Mijnbouw Genootschap Nederland en Koloniën, Geology Series*, vol. 8, p. 617–631, pls. 1–4.
- Yabe, H. and Hanzawa, S., 1929: Tertiary foraminiferous rocks of the Philippines. *Science Reports of the Tohoku University, Second Series (Geology)*, vol. 11, no. 3, p. 137–190, pls. 15–27.